Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.



enro Enro

ET-34



December 1934

A LABORATORY DEVICE FOR SUPPLYING LIQUIDS TO ADULT INSECTS

By Heber C. Donohoe, Division of Fruit Insects, Bureau of Entomology and Plant Quarantine, U. S. Department of Agriculture

In the writer's life-history studies of the raisin moth, <u>Ephestia figulilella</u> Gregson, in the dried-fruit insect laboratory at Fresno, Calif., it was early noted that the number of eggs laid per female was increased and the length of life of both sexes prolonged if the adults used were supplied with either water or sugar solution. The major difficulty encountered in attempting to supply this need was the fact that, when various methods of affording liquid food were used, the moths became entangled in the fluid; then, in attempting to fly about their containers, their wings stuck to the glass or celluloid and they shortly died.

The drinking cup finally devised eliminates the possibility of the insects themselves getting wet, yet provides a readily available supply of liquid. This cup is fashioned from a section of $\frac{1}{8}$ —inch bore glass tubing, $\frac{3}{8}$ inch long, one end of which is fused and rounded over a flame, thus forming a miniature receptacle. When in use, the cup is inserted in a large raisin so that the open end is flush with the surface. With the mouth of the cup upward, it may be filled with an ordinary medicine dropper, the glass of which is drawn out into a capillary tube.

The liquid cannot be spilled out of the cup since, in a tube of such small size, the interfacial tension of glass and fluid is sufficient to prevent it. A moth will feed readily from such a container by thrusting its proboscis into the opening of the cup; yet, owing to the small size of this opening, its body is kept entirely free of the liquid within.



SSCS 100 Standill

A LABORATORY DEVICE FOR SUPPLYING LIGHTLE TO ADULT DESCRIP

By Heber C. Donohoe, Division of Fruit Inmedia, Europa of Entomology and Plant Quarantine, U. S. Dopartment of Agriculture

In the writer's life-history studies of his mists acts, financial installa Gregson, in the dried-front insent laboratory as fractor if was early noted that the number of ears laid per leadle win resead and the longth of life of boils same prolonged if the adults deep mility encountered in attempting to simply this need was use fact that a various methods of affording liquid food ware-wred, the molbs because antangled in the fluid; then, in attempting to fly shoul light desirable dand they should the claimers, their wings attent to the glass or cellulated and they shortly

The drinking cup finally dayted stainance the possibility of the sects themselves getting wet, yet provides a concile available equity liquid. This cup is leshioned from a section of 1-inch bore glass sine, I know long, one and of which is fused and rounded over a fidne, we forming a minister receptable. When in use, the cup if the cup upward, it may be filled with the rather life and the glass of which is drawn out into a capillary taken

The liquid cannot be spilled out of the cup sides in a twhe of the small size, the interfacial tension of glass and Theid is surficient to prevent it. A noth will feed readily from such a container be liquid; its probosels into the opening of the cup; yet, owing to the small se of this opening, its body is kept entirely free of the liquid within.